

STIELTJES LIKE FUNCTIONS AND INVERSE PROBLEMS FOR SYSTEMS WITH SCHRÖDINGER OPERATOR

SERGEY BELYI AND EDUARD TSEKANOVSKII

ABSTRACT. A class of scalar Stieltjes like functions is realized as linear-fractional transformations of transfer functions of conservative systems based on a Schrödinger operator T_h in $L_2[a, +\infty)$ with a non-selfadjoint boundary condition. In particular it is shown that any Stieltjes function of this class can be realized in the unique way so that the main operator $\mathbb{A}$ of a system is an accretive $(*)$ -extension of a Schrödinger operator T_h . We derive formulas that restore the system uniquely and allow to find the exact value of a non-real parameter h in the definition of T_h as well as a real parameter μ that appears in the construction of the elements of the realizing system. An elaborate investigation of these formulas shows the dynamics of the restored parameters h and μ in terms of the changing free term γ from the integral representation of the realizable function. It turns out that the parametric equations for the restored parameter h represent different circles whose centers and radii are determined by the realizable function. Similarly, the behavior of the restored parameter μ are described by hyperbolas.

1. INTRODUCTION

Realizations of different classes of holomorphic operator-valued functions in the open right half-plane, unit circle, and upper half-plane, as well as inverse spectral problems, play an important role in the spectral analysis of non-self-adjoint operators, interpolation problems, and system theory. The literature on realization theory is too extensive to be discussed exhaustively in this note. We refer, however, to [2], [3], [7], [8], [9], [11], [12], [18], [24], [28] and the literature therein. A class of Herglotz-Nevanlinna functions is a rich source for many types of realization problems. An operator-valued function $V(z)$ acting on a finite-dimensional Hilbert space E belongs to the class of operator-valued Herglotz-Nevanlinna functions if it is holomorphic on $\mathbb{C} \setminus \mathbb{R}$, if it is symmetric with respect to the real axis, i.e., $V(z)^* = V(\bar{z})$, $z \in \mathbb{C} \setminus \mathbb{R}$, and if it satisfies the positivity condition

$$\operatorname{Im} V(z) \geq 0, \quad z \in \mathbb{C}_+.$$

It is well known (see e.g. [16], [17]) that operator-valued Herglotz-Nevanlinna functions admit the following integral representation:

$$(1.1) \quad V(z) = Q + Lz + \int_{\mathbb{R}} \left(\frac{1}{t-z} - \frac{t}{1+t^2} \right) dG(t), \quad z \in \mathbb{C} \setminus \mathbb{R},$$

Date: February 1, 2008.

2000 Mathematics Subject Classification. Primary 47A10, 47B44; Secondary 46E20, 46F05.

Key words and phrases. Operator colligation, conservative and impedance system, transfer (characteristic) function.

where $Q = Q^*$, $L \geq 0$, and $G(t)$ is a nondecreasing operator-valued function on $\mathbb{R}$ with values in the class of nonnegative operators in E such that

$$(1.2) \quad \int_{\mathbb{R}} \frac{(dG(t)x, x)_E}{1+t^2} < \infty, \quad x \in E.$$

The realization of a selected class of Herglotz-Nevanlinna functions is provided by a linear conservative system Θ of the form

$$(1.3) \quad \begin{cases} (\mathbb{A} - zI)x = KJ\varphi_- \\ \varphi_+ = \varphi_- - 2iK^*x \end{cases}$$

or

$$(1.4) \quad \Theta = \begin{pmatrix} \mathbb{A} & K & J \\ \mathcal{H}_+ \subset \mathcal{H} \subset \mathcal{H}_- & & E \end{pmatrix}.$$

In this system $\mathbb{A}$, the *main operator* of the system, is a so-called $(*)$ -extension, which is a bounded linear operator from $\mathcal{H}_+$ into $\mathcal{H}_-$, where $\mathcal{H}_+ \subset \mathcal{H} \subset \mathcal{H}_-$ is a rigged Hilbert space. Moreover, K is a bounded linear operator from the finite-dimensional Hilbert space E into $\mathcal{H}_-$, while $J = J^* = J^{-1}$ is acting on E , are such that $\text{Im } \mathbb{A} = KJK^*$. Also, $\varphi_- \in E$ is an input vector, $\varphi_+ \in E$ is an output vector, and $x \in \mathcal{H}_+$ is a vector of the state space of the system Θ . The system described by (1.3)-(1.4) is called a rigged canonical system of the Livšic type [22] or the Brodskiĭ-Livšic rigged operator colligation, cf., e.g. [11], [12], [13]. The operator-valued function

$$(1.5) \quad W_{\Theta}(z) = I - 2iK^*(\mathbb{A} - zI)^{-1}KJ$$

is a transfer function (or characteristic function) of the system Θ . It was shown in [11] that an operator-valued function $V(z)$ acting on a Hilbert space E of the form (1.1) can be represented and realized in the form

$$(1.6) \quad V(z) = i[W_{\Theta}(z) + I]^{-1}[W_{\Theta}(z) - I] = K^*(\mathbb{A}_R - zI)^{-1}K,$$

where $W_{\Theta}(z)$ is a transfer function of some canonical scattering ($J = I$) system Θ , and where the “*real part*” $\mathbb{A}_R = \frac{1}{2}(\mathbb{A} + \mathbb{A}^*)$ of $\mathbb{A}$ satisfies $\mathbb{A}_R \supset A$ if and only if the function $V(z)$ in (1.1) satisfies the following two conditions:

$$(1.7) \quad \begin{cases} L = 0, \\ Qx = \int_{\mathbb{R}} \frac{t}{1+t^2} dG(t)x \quad \text{when} \quad \int_{\mathbb{R}} (dG(t)x, x)_E < \infty. \end{cases}$$

In the current paper we are going to focus on an important subclass of Herglotz-Nevanlinna functions, the so called Stieltjes like functions that also includes Stieltjes functions. In Section 4 we specify a subclass of realizable Stieltjes operator-functions and show that any member of this subclass can be realized by a system of the form (1.4) whose main operator $\mathbb{A}$ is accretive.

In Section 5 we introduce a class of Stieltjes like scalar functions. Then we rely on the general realization results developed in Section 4 (see also [15]) to restore a system Θ of the form (1.4) containing the Schrödinger operator in $L_2[a, +\infty)$ with non-self-adjoint boundary conditions

$$\begin{cases} T_h y = -y'' + q(x)y \\ y'(a) = h y(a) \end{cases}, \quad \left(q(x) = \overline{q(x)}, \text{Im } h \neq 0 \right).$$

We show that if a non-decreasing function $\sigma(t)$ is the spectral distribution function of positive self-adjoint boundary value problem

$$\begin{cases} A_\theta y = -y'' + q(x)y \\ y'(a) = \theta y(a) \end{cases}$$

and satisfies conditions

$$\int_0^\infty d\sigma(t) = \infty, \quad \int_0^\infty \frac{d\sigma(t)}{1+t} < \infty,$$

then for every real γ a Stieltjes like function

$$V(z) = \gamma + \int_0^\infty \frac{d\sigma(t)}{t-z}$$

can be realized in the unique way as a $V_\Theta(z)$ function of a rigged canonical system Θ containing some Schrödinger operator T_h . In particular, it is shown that for every $\gamma \geq 0$ a Stieltjes function $V(z)$ with integral representation above can be realized by a system Θ whose main operator $\mathbb{A}$ is an accretive $(*)$ -extension of a Schrödinger operator T_h .

On top of the general realization results, Section 5 provides the reader with formulas that allow to find the exact value of a non-real parameter h in the definition of T_h of the realizing system Θ . Similar investigation is presented in Section 6 to describe the real parameter μ that appears in the construction of the elements of the realizing system. A detailed study of these formulas shows the dynamics of the restored parameters h and μ in terms of a changing free term γ in the integral representation of $V(z)$ above. It will be shown and graphically presented that the parametric equations for the restored parameter h represent different circles whose centers and radii are completely determined by the function $V(z)$. Similarly, the behavior of the restored parameter μ are described by hyperbolas.

2. SOME PRELIMINARIES

For a pair of Hilbert spaces $\mathcal{H}_1, \mathcal{H}_2$ we denote by $[\mathcal{H}_1, \mathcal{H}_2]$ the set of all bounded linear operators from $\mathcal{H}_1$ to $\mathcal{H}_2$. Let A be a closed, densely defined, symmetric operator in a Hilbert space $\mathcal{H}$ with inner product $(f, g), f, g \in \mathcal{H}$. Consider the rigged Hilbert space

$$\mathcal{H}_+ \subset \mathcal{H} \subset \mathcal{H}_-,$$

where $\mathcal{H}_+ = D(A^*)$ and

$$(f, g)_+ = (f, g) + (A^*f, A^*g), \quad f, g \in D(A^*).$$

Note that identifying the space conjugate to $\mathcal{H}_\pm$ with $\mathcal{H}_\mp$, we get that if $\mathbb{A} \in [\mathcal{H}_+, \mathcal{H}_-]$ then $\mathbb{A}^* \in [\mathcal{H}_+, \mathcal{H}_-]$.

Definition 2.1. An operator $\mathbb{A} \in [\mathcal{H}_+, \mathcal{H}_-]$ is called a self-adjoint bi-extension of a symmetric operator A if $\mathbb{A} = \mathbb{A}^*$, $\mathbb{A} \supset A$, and the operator

$$\widehat{A}f = \mathbb{A}f, \quad f \in D(\widehat{A}) = \{f \in \mathcal{H}_+ : \mathbb{A}f \in \mathcal{H}\}$$

is self-adjoint in $\mathcal{H}$.

The operator $\widehat{A}$ in the above definition is called a *quasi-kernel* of a self-adjoint bi-extension $\mathbb{A}$ (see [27]).

Definition 2.2. An operator $\mathbb{A} \in [\mathcal{H}_+, \mathcal{H}_-]$ is called a $(*)$ -extension (or correct bi-extension) of an operator T (with non-empty set $\rho(T)$ of regular points) if

$$\mathbb{A} \supset T \supset A, \mathbb{A}^* \supset T^* \supset A$$

and the operator $\mathbb{A}_R = \frac{1}{2}(\mathbb{A} + \mathbb{A}^*)$ is a self-adjoint bi-extension of an operator A .

The existence, description, and analog of von Neumann's formulas for self-adjoint bi-extensions and $(*)$ -extensions were discussed in [27] (see also [4], [5], [11]). For instance, if Φ is an isometric operator from the defect subspace $\mathfrak{N}_i$ of the symmetric operator A onto the defect subspace $\mathfrak{N}_{-i}$, then the formulas below establish a one-to one correspondence between $(*)$ -extensions of an operator T and Φ

$$(2.1) \quad \mathbb{A}f = A^*f + iR(\Phi - I)x, \quad \mathbb{A}^*f = A^*f + iR(\Phi - I)y,$$

where $x, y \in \mathfrak{N}_i$ are uniquely determined from the conditions

$$f - (\Phi + I)x \in D(T), \quad f - (\Phi + I)y \in D(T^*)$$

and R is the Riesz-Berezanskii operator of the triplet $\mathcal{H}_+ \subset \mathcal{H} \subset \mathcal{H}_-$ that maps $\mathcal{H}_+$ isometrically onto $\mathcal{H}_-$ (see [27]). If the symmetric operator A has deficiency indices (n, n) , then formulas (2.1) can be rewritten in the following form

$$(2.2) \quad \mathbb{A}f = A^*f + \sum_{k=1}^n \Delta_k(f)V_k, \quad \mathbb{A}^*f = A^*f + \sum_{k=1}^n \delta_k(f)V_k,$$

where $\{V_j\}_1^n \in \mathcal{H}_-$ is a basis in the subspace $R(\Phi - I)\mathfrak{N}_i$, and $\{\Delta_k\}_1^n, \{\delta_k\}_1^n$ are bounded linear functionals on $\mathcal{H}_+$ with the properties

$$(2.3) \quad \Delta_k(f) = 0, \quad \forall f \in D(T), \quad \delta_k(f) = 0, \quad \forall f \in D(T^*).$$

Let $\mathcal{H} = L_2[a, +\infty)$ and $l(y) = -y'' + q(x)y$ where q is a real locally summable function. Suppose that the symmetric operator

$$(2.4) \quad \begin{cases} Ay = -y'' + q(x)y \\ y(a) = y'(a) = 0 \end{cases}$$

has deficiency indices $(1, 1)$. Let D^* be the set of functions locally absolutely continuous together with their first derivatives such that $l(y) \in L_2[a, +\infty)$. Consider $\mathcal{H}_+ = D(A^*) = D^*$ with the scalar product

$$(y, z)_+ = \int_a^\infty \left(y(x)\overline{z(x)} + l(y)\overline{l(z)} \right) dx, \quad y, z \in D^*.$$

Let

$$\mathcal{H}_+ \subset L_2[a, +\infty) \subset \mathcal{H}_-$$

be the corresponding triplet of Hilbert spaces. Consider operators

$$(2.5) \quad \begin{cases} T_h y = l(y) = -y'' + q(x)y \\ h y(a) - y'(a) = 0 \end{cases}, \quad \begin{cases} T_h^* y = l(y) = -y'' + q(x)y \\ \overline{h} y(a) - y'(a) = 0 \end{cases},$$

$$\begin{cases} \widehat{A} y = l(y) = -y'' + q(x)y \\ \mu y(a) - y'(a) = 0 \end{cases}, \quad \text{Im } \mu = 0.$$

It is well known [1] that $\widehat{A} = \widehat{A}^*$. The following theorem was proved in [6].

Theorem 2.3. *The set of all $(*)$ -extensions of a non-self-adjoint Schrödinger operator T_h of the form (2.5) in $L_2[a, +\infty)$ can be represented in the form*

$$(2.6) \quad \begin{aligned} \mathbb{A}y &= -y'' + q(x)y - \frac{1}{\mu - h} [y'(a) - hy(a)] [\mu\delta(x-a) + \delta'(x-a)], \\ \mathbb{A}^*y &= -y'' + q(x)y - \frac{1}{\mu - \bar{h}} [y'(a) - \bar{h}y(a)] [\mu\delta(x-a) + \delta'(x-a)]. \end{aligned}$$

In addition, the formulas (2.6) establish a one-to-one correspondence between the set of all $(*)$ -extensions of a Schrödinger operator T_h of the form (2.5) and all real numbers $\mu \in [-\infty, +\infty]$.

Definition 2.4. An operator T with the domain $D(T)$ and $\rho(T) \neq \emptyset$ acting on a Hilbert space $\mathcal{H}$ is called *accretive* if

$$\operatorname{Re}(Tf, f) \geq 0, \quad \forall f \in D(T).$$

Definition 2.5. An accretive operator T is called [20] α -sectorial if there exists a value of $\alpha \in (0, \pi/2)$ such that

$$\cot \alpha |\operatorname{Im}(Tf, f)| \leq \operatorname{Re}(Tf, f), \quad f \in \mathcal{D}(T).$$

An accretive operator is called *extremal accretive* if it is not α -sectorial for any $\alpha \in (0, \pi/2)$.

Consider the symmetric operator A of the form (2.4) with defect indices (1,1), generated by the differential operation $l(y) = -y'' + q(x)y$. Let $\varphi_k(x, \lambda)$ ($k = 1, 2$) be the solutions of the following Cauchy problems:

$$\begin{cases} l(\varphi_1) = \lambda\varphi_1 \\ \varphi_1(a, \lambda) = 0 \\ \varphi_1'(a, \lambda) = 1 \end{cases}, \quad \begin{cases} l(\varphi_2) = \lambda\varphi_2 \\ \varphi_2(a, \lambda) = -1 \\ \varphi_2'(a, \lambda) = 0 \end{cases}.$$

It is well known [1] that there exists a function $m_\infty(\lambda)$ (called the Weyl-Titchmarsh function) for which

$$\varphi(x, \lambda) = \varphi_2(x, \lambda) + m_\infty(\lambda)\varphi_1(x, \lambda)$$

belongs to $L_2[a, +\infty)$.

Suppose that the symmetric operator A of the form (2.4) with deficiency indices (1,1) is nonnegative, i.e., $(Af, f) \geq 0$ for all $f \in D(A)$. It was shown in [25] that the Schrödinger operator T_h of the form (2.5) is accretive if and only if

$$(2.7) \quad \operatorname{Re} h \geq -m_\infty(-0).$$

For real h such that $h \geq -m_\infty(-0)$ we get a description of all nonnegative self-adjoint extensions of an operator A . For $h = -m_\infty(-0)$ the corresponding operator

$$(2.8) \quad \begin{cases} A_K y = -y'' + q(x)y \\ y'(a) + m_\infty(-0)y(a) = 0 \end{cases}$$

is the Kreĭn-von Neumann extension of A and for $h = +\infty$ the corresponding operator

$$(2.9) \quad \begin{cases} A_F y = -y'' + q(x)y \\ y(a) = 0 \end{cases}$$

is the Friedrichs extension of A (see [25], [6]).

3. RIGGED CANONICAL SYSTEMS WITH SCHRÖDINGER OPERATOR

Let $\mathbb{A}$ be $(*)$ - extension of an operator T , i.e.,

$$\mathbb{A} \supset T \supset A, \quad \mathbb{A}^* \supset T^* \supset A$$

where A is a symmetric operator with deficiency indices (n, n) and $D(A) = D(T) \cap D(T^*)$. In what follows we will only consider the case when the symmetric operator A has dense domain, i.e., $\overline{\mathcal{D}(A)} = \mathcal{H}$.

Definition 3.1. A system of equations

$$\begin{cases} (\mathbb{A} - zI)x = KJ\varphi_- \\ \varphi_+ = \varphi_- - 2iK^*x \end{cases},$$

or an array

$$(3.1) \quad \Theta = \begin{pmatrix} \mathbb{A} & K & J \\ \mathcal{H}_+ \subset \mathcal{H} \subset \mathcal{H}_- & & E \end{pmatrix}$$

is called a *rigged canonical system of the Livsic type* or the *Brodskiĭ-Livsic rigged operator colligation* if:

- 1) E is a finite-dimensional Hilbert space with scalar product $(\cdot, \cdot)_E$ and the operator J in this space satisfies the conditions $J = J^* = J^{-1}$,
- 2) $K \in [E, \mathcal{H}_-]$,
- 3) $\text{Im } \mathbb{A} = KJK^*$, where $K^* \in [\mathcal{H}_+, E]$ is the adjoint of K .

In the definition above $\varphi_- \in E$ stands for an input vector, $\varphi_+ \in E$ is an output vector, and x is a state space vector in $\mathcal{H}$. An operator $\mathbb{A}$ is called a *main operator* of the system Θ , J is a *direction operator*, and K is a *channel operator*. An operator-valued function

$$(3.2) \quad W_\Theta(\lambda) = I - 2iK^*(\mathbb{A} - \lambda I)^{-1}KJ$$

defined on the set $\rho(T)$ of regular points of an operator T is called the *transfer function* (*characteristic function*) of the system Θ , i.e., $\varphi_+ = W_\Theta(\lambda)\varphi_-$. It is known [25], [27] that any $(*)$ -extension $\mathbb{A}$ of an operator T ($\mathbb{A}^* \supset T \supset A$), where A is a symmetric operator with deficiency indices (n, n) ($n < \infty$), $D(A) = D(T) \cap D(T^*)$, can be included as a main operator of some rigged canonical system with $\dim E < \infty$ and invertible channel operator K .

It was also established [25], [27] that

$$(3.3) \quad V_\Theta(\lambda) = K^*(\text{Re } \mathbb{A} - \lambda I)^{-1}K$$

is a Herglotz-Nevanlinna operator-valued function acting on a Hilbert space E , satisfying the following relation for $\lambda \in \rho(T)$, $\text{Im } \lambda \neq 0$

$$(3.4) \quad V_\Theta(\lambda) = i[W_\Theta(\lambda) - I][W_\Theta(\lambda) + I]^{-1}J.$$

Alternatively,

$$(3.5) \quad \begin{aligned} W_\Theta(\lambda) &= (I + iV_\Theta(\lambda)J)^{-1}(I - iV_\Theta(\lambda)J) \\ &= (I - iV_\Theta(\lambda)J)(I + iV_\Theta(\lambda)J)^{-1}. \end{aligned}$$

Let us recall (see [27], [6]) that a symmetric operator with dense domain $\mathcal{D}(A)$ is called *prime* if there is no reducing, nontrivial invariant subspace on which A

induces a self-adjoint operator. It was established in [26] that a symmetric operator A is prime if and only if

$$(3.6) \quad \text{c.l.s.}_{\lambda \neq \bar{\lambda}} \mathfrak{N}_\lambda = \mathcal{H}.$$

We call a rigged canonical system of the form (3.1) *prime* if

$$\text{c.l.s.}_{\lambda \neq \bar{\lambda}, \lambda \in \rho(T)} \mathfrak{N}_\lambda = \mathcal{H}.$$

One easily verifies that if system Θ is prime, then a symmetric operator A of the system is prime as well.

The following theorem [6] establishes the connection between two rigged canonical systems with equal transfer functions.

Theorem 3.2. *Let $\Theta_1 = \begin{pmatrix} \mathbb{A}_1 & K_1 & J \\ \mathcal{H}_{+1} \subset \mathcal{H}_1 \subset \mathcal{H}_{-1} & & E \end{pmatrix}$ and $\Theta_2 = \begin{pmatrix} \mathbb{A}_2 & K_2 & J \\ \mathcal{H}_{+2} \subset \mathcal{H}_2 \subset \mathcal{H}_{-2} & & E \end{pmatrix}$ be two prime rigged canonical systems of the Livsic type with*

$$(3.7) \quad \begin{aligned} \mathbb{A}_1 \supset T_1 \supset A_1, \quad \mathbb{A}_1^* \supset T_1^* \supset A_1, \\ \mathbb{A}_2 \supset T_2 \supset A_2, \quad \mathbb{A}_2^* \supset T_2^* \supset A_2, \end{aligned}$$

and such that A_1 and A_2 have finite and equal defect indices.

If

$$(3.8) \quad W_{\Theta_1}(\lambda) = W_{\Theta_2}(\lambda),$$

then there exists an isometric operator U from $\mathcal{H}_1$ onto $\mathcal{H}_2$ such that $U_+ = U|_{\mathcal{H}_{+1}}$ is an isometry¹ from $\mathcal{H}_{+1}$ onto $\mathcal{H}_{+2}$, $U_-^* = U_+^*$ is an isometry from $\mathcal{H}_{-1}$ onto $\mathcal{H}_{-2}$, and

$$(3.9) \quad UT_1 = T_2U, \quad \mathbb{A}_2 = U_- \mathbb{A}_1 U_+^{-1}, \quad U_- K_1 = K_2.$$

Corollary 3.3. *Let Θ_1 and Θ_2 be the two prime systems from the statement of theorem 3.2. Then the mapping U described in the conclusion of the theorem is unique.*

Proof. First let us make an observation that if $\Theta = \begin{pmatrix} \mathbb{A} & K & J \\ \mathcal{H}_+ \subset \mathcal{H} \subset \mathcal{H}_- & & E \end{pmatrix}$ is a prime rigged canonical system such that $U_- \mathbb{A} = \mathbb{A} U_+$ and $U_- K = K$, where U is an isometry mapping described in theorem 3.2, then $U = I$. Indeed, it is well known [27] that

$$(3.10) \quad (\text{Re } \mathbb{A} - \lambda I)^{-1} K E = \mathfrak{N}_\lambda.$$

We have

$$\begin{aligned} U(\text{Re } \mathbb{A} - \lambda I)^{-1} K e &= U_+(\text{Re } \mathbb{A} - \lambda I)^{-1} K e = (\text{Re } \mathbb{A} - \lambda I)^{-1} U_- K e \\ &= (\text{Re } \mathbb{A} - \lambda I)^{-1} K e, \quad \forall e \in E, \lambda \neq \bar{\lambda}. \end{aligned}$$

Combining the above equation with (3.6) and (3.10) we obtain $U = I$.

¹It was shown in [6] that the operator U_+ defined this way is an isometry from $\mathcal{H}_{+1}$ onto $\mathcal{H}_{+2}$. It is also shown there that the isometric operator $U^* : \mathcal{H}_{+2} \rightarrow \mathcal{H}_{+1}$ uniquely defines operator $U_- = (U^*)^* : \mathcal{H}_{-1} \rightarrow \mathcal{H}_{-2}$.

Now let Θ_1 and Θ_2 be the two prime systems from the statement of theorem 3.2. Suppose there are two isometric mappings U_1 and U_2 guaranteed by theorem 3.2. Then the relations

$$\mathbb{A}_2 = U_{-,1}\mathbb{A}_1U_{+,1}^{-1}, \quad U_{-,1}K_1 = K_2, \quad \mathbb{A}_2 = U_{-,2}\mathbb{A}_1U_{+,2}^{-1}, \quad U_{-,2}K_1 = K_2,$$

lead to

$$\mathbb{A}_1U_{+,1}^{-1}U_{+,2} = U_{-,1}^{-1}U_{-,2}\mathbb{A}_1, \quad U_{-,1}^{-1}U_{-,2}K = K.$$

Since Θ_1 is prime then $U_1^{-1}U_2 = I$ and hence $U_1 = U_2$. This proves the uniqueness of U . $\square$

Now we shall construct a rigged canonical system based on a non-self-adjoint Schrödinger operator. One can easily check that the $(*)$ -extension

$$\mathbb{A}y = -y'' + q(x)y - \frac{1}{\mu - h} [y'(a) - hy(a)] [\mu\delta(x-a) + \delta'(x-a)], \quad \text{Im } h > 0$$

of the non-self-adjoint Schrödinger operator T_h of the form (2.5) satisfies the condition

$$(3.11) \quad \text{Im } \mathbb{A} = \frac{\mathbb{A} - \mathbb{A}^*}{2i} = (\cdot, g)g,$$

where

$$(3.12) \quad g = \frac{(\text{Im } h)^{\frac{1}{2}}}{|\mu - h|} [\mu\delta(x-a) + \delta'(x-a)]$$

and $\delta(x-a)$, $\delta'(x)$ are the delta-function and its derivative at the point a . Moreover,

$$(3.13) \quad (y, g) = \frac{(\text{Im } h)^{\frac{1}{2}}}{|\mu - h|} [\mu y(a) - y'(a)],$$

where

$$y \in \mathcal{H}_+, g \in \mathcal{H}_-, \mathcal{H}_+ \subset L_2(a, +\infty) \subset \mathcal{H}_-$$

and the triplet of Hilbert spaces is as discussed in theorem 2.3. Let $E = \mathbb{C}$, $Kc = cg$ ($c \in \mathbb{C}$). It is clear that

$$(3.14) \quad K^*y = (y, g), \quad y \in \mathcal{H}_+$$

and $\text{Im } \mathbb{A} = KK^*$. Therefore, the array

$$(3.15) \quad \Theta = \begin{pmatrix} \mathbb{A} & K & 1 \\ \mathcal{H}_+ \subset L_2[a, +\infty) \subset \mathcal{H}_- & \mathbb{C} & \end{pmatrix}$$

is a rigged canonical system with the main operator $\mathbb{A}$ of the form (2.6), the direction operator $J = 1$ and the channel operator K of the form (3.14). Our next logical step is finding the transfer function of (3.15). It was shown in [6] that

$$(3.16) \quad W_\Theta(\lambda) = \frac{\mu - h}{\mu - \bar{h}} \frac{m_\infty(\lambda) + \bar{h}}{m_\infty(\lambda) + h},$$

and

$$(3.17) \quad V_\Theta(\lambda) = \frac{(m_\infty(\lambda) + \mu) \text{Im } h}{(\mu - \text{Re } h) m_\infty(\lambda) + \mu \text{Re } h - |h|^2}.$$

4. REALIZATION OF STIELTJES FUNCTIONS

Let E be a finite-dimensional Hilbert space. The scalar versions of the following definition can be found in [19].

Definition 4.1. We will call an operator-valued Herglotz-Nevanlinna function $V(z) \in [E, E]$ by a *Stieltjes function* if $V(z)$ admits the following integral representation

$$(4.1) \quad V(z) = \gamma + \int_0^\infty \frac{dG(t)}{t - z},$$

where $\gamma \geq 0$ and $G(t)$ is a non-decreasing on $[0, +\infty)$ operator-valued function such that

$$\int_0^\infty \frac{(dG(t)e, e)_E}{1+t} < \infty, \quad \forall e \in E.$$

Alternatively (see [19]) an operator-valued function $V(z)$ is Stieltjes if it is holomorphic in $\text{Ext}[0, +\infty)$ and

$$(4.2) \quad \frac{\text{Im}[zV(z)]}{\text{Im} z} \geq 0.$$

The theorem 4.2 below was stated in [14], [15] and we present its proof for the convenience of a reader.

Theorem 4.2. *Let Θ be a prime system of the form (3.1). Then an operator-valued function $V_\Theta(z)$ defined by (3.3), (3.4) is a Stieltjes function if and only if the main operator $\mathbb{A}$ of the system Θ is accretive.*

Proof. Let us assume first that $\mathbb{A}$ is an accretive operator, i.e. $(\text{Re } \mathbb{A}x, x) \geq 0$, for all $x \in \mathcal{H}_+$. Let $\{z_k\}$ ($k = 1, \dots, n$) be a sequence of non-real complex numbers and h_k be a sequence of vectors in E . Let us denote

$$(4.3) \quad Kh_k = \delta_k, \quad x_k = (\text{Re } \mathbb{A} - z_k I)^{-1} \delta_k, \quad x = \sum_{k=1}^n x_k.$$

Since $(\text{Re } \mathbb{A}x, x) \geq 0$, we have

$$(4.4) \quad \sum_{k,l=1}^n (\text{Re } \mathbb{A}x_k, x_l) \geq 0.$$

By formal calculations one can have

$$\text{Re } \mathbb{A}x_k = \delta_k + z_k(\text{Re } \mathbb{A} - z_k I)^{-1} \delta_k,$$

and

$$\begin{aligned} \sum_{k,l=1}^n (\text{Re } \mathbb{A}x_k, x_l) &= \sum_{k,l=1}^n [(\delta_k, (\text{Re } \mathbb{A} - z_l I)^{-1} \delta_l) \\ &\quad + (z_k(\text{Re } \mathbb{A} - z_k I)^{-1} \delta_k, (\text{Re } \mathbb{A} - z_l I)^{-1} \delta_l)]. \end{aligned}$$

Using obvious equalities

$$((\text{Re } \mathbb{A} - z_k I)^{-1} Kh_k, Kh_l) = (V_\Theta(z_k)h_k, h_l)_E,$$

and

$$((\operatorname{Re} \mathbb{A} - \bar{z}_l I)^{-1} (\operatorname{Re} \mathbb{A} - z_k I)^{-1} K h_k, K h_l) = \left(\frac{V_\theta(z_k) - V_\theta(\bar{z}_l)}{z_k - \bar{z}_l} h_k, h_l \right)_E,$$

we obtain

$$(4.5) \quad \sum_{k,l=1}^n (\operatorname{Re} \mathbb{A} x_k, x_l) = \sum_{k,l=1}^n \left(\frac{z_k V_\theta(z_k) - \bar{z}_l V_\theta(\bar{z}_l)}{z_k - \bar{z}_l} h_k, h_l \right)_E \geq 0.$$

The choice of z_k was arbitrary, which means that $V_\Theta(z)$ is a Stieltjes function (see [3]).

Now we prove necessity. Since Θ is a prime system then A is a prime symmetric operator. Then the equivalence of (4.5) and (4.4) implies that $(\operatorname{Re} \mathbb{A} x, x) \geq 0$ for any x from c.l.s. $\{\mathfrak{N}_z\}$, $z \neq \bar{z}$. As we have already mentioned above, a symmetric operator A with the equal deficiency indices is prime if and only if for all $\lambda \neq \bar{\lambda}$

$$\text{c.l.s. } \{\mathfrak{N}_\lambda\} = \mathcal{H}.$$

Therefore we can conclude that $(\operatorname{Re} \mathbb{A} x, x) \geq 0$ for any $x \in \mathcal{H}_+$ and hence $\mathbb{A}$ is an accretive operator. $\square$

A system Θ of the form (3.1) is called an *accretive system* if its main operator $\mathbb{A}$ is accretive.

Now we define a certain class $S_0(R)$ of realizable Stieltjes functions. At this point we need to note that since Stieltjes functions form a subset of Herglotz-Nevanlinna functions then we can utilize the conditions (1.7) to form a *class* $S(R)$ of all *realizable Stieltjes functions* (see also [15]). Clearly, $S(R)$ is a subclass of $N(R)$ of all realizable Herglotz-Nevanlinna functions described in details in [11] and [12]. To see the specifications of the class $S(R)$ we recall that aside of integral representation (4.1), any Stieltjes function admits a representation (1.1). Applying condition (1.7) we obtain

$$(4.6) \quad Q = \frac{1}{2} [V_\theta(-i) + V_\theta^*(-i)] = \gamma + \int_0^{+\infty} \frac{t}{1+t^2} dG(t).$$

Combining the second part of condition (1.7) and (4.6) we conclude that

$$(4.7) \quad \gamma e = 0,$$

for all $e \in E$ such that

$$(4.8) \quad \int_0^\infty (dG(t)e, e)_E < \infty.$$

holds. Consequently, (4.7)-(4.8) is precisely the condition for $V(z) \in S(R)$.

We are going to focus though on the subclass $S_0(R)$ of $S(R)$ whose definition is the following.

Definition 4.3. An operator-valued Stieltjes function $V(z) \in [E, E]$ is said to be a member of the **class** $S_0(R)$ if in the representation (4.1) we have

$$(4.9) \quad \int_0^\infty (dG(t)e, e)_E = \infty.$$

for all non-zero $e \in E$.

We note that a function $V(z)$ can belong to class $S_0(R)$ and have an arbitrary constant $\gamma \geq 0$ in the representation (4.1).

The following statement [15] is the direct realization theorem for the functions of the class $S_0(R)$.

Theorem 4.4. *Let Θ be an accretive system of the form (3.1). Then the operator-function $V_\Theta(z)$ of the form (3.3), (3.4) belongs to the class $S_0(R)$.*

Proof. To see that $V_\Theta(z)$ is a Stieltjes operator-function we merely apply theorem 4.2 to system Θ .

Now we will show that $V_\Theta(z)$ belongs to $S_0(R)$. It was shown in [11] and [12] that $E_\infty = K^{-1}\mathfrak{L}$, where $\mathfrak{L} = \mathcal{H} \ominus \overline{\mathcal{D}(A)}$ and

$$E_\infty = \left\{ e \in E : \int_0^{+\infty} (dG(t)e, e)_E < \infty \right\}.$$

But $\overline{\mathcal{D}(A)} = \mathcal{H}$ and consequently $\mathfrak{L} = \{0\}$. Next, $E_\infty = \{0\}$,

$$\int_0^\infty (dG(t)e, e)_E = \infty,$$

for all non-zero $e \in E$, and therefore $V_\theta(z) \in S_0(R)$. □

The inverse realization theorem can be stated and proved (see [15]) for the classes $S_0(R)$ as follows.

Theorem 4.5. *Let a operator-valued function $V(z)$ belong to the class $S_0(R)$. Then $V(z)$ admits a realization by an accretive prime system Θ of the form (3.1) with $\mathcal{D}(T) \neq \mathcal{D}(T^*)$ and $J = I$.*

Proof. We have already noted that the class of Stieltjes function lies inside the wider class of all Herglotz-Nevanlinna functions. Thus all we actually have to show is that $S_0(R) \subset N_0(R)$, where $N_0(R)$ is subclass of realizable Herglotz-Nevanlinna functions described in [12], and that the realizing system constructed in [12] appears to be an accretive system. The former is rather obvious and follows directly from the definition of the class $S_0(R)$. To see that the realizing system is accretive we need to apply theorem 4.2 to $V_\theta(z) = V(z)$, where $V_\theta(z)$ is related to the model system Θ that was constructed in [12]. As it was also shown in [11] and [12], the symmetric operator A of the model system Θ is prime and hence (3.6) takes place. We are going to show that in this case the system Θ is also prime, i.e.,

$$(4.10) \quad \text{c.l.s.} \quad \mathfrak{N}_\lambda = \mathcal{H}.$$

$\lambda \neq \bar{\lambda}, \lambda \in \rho(T)$

Consider the operator $U_{\lambda_0\lambda} = (\tilde{A} - \lambda_0 I)(\tilde{A} - \lambda I)^{-1}$, where $\tilde{A}$ is an arbitrary self-adjoint extension of A . By a simple check one confirms that $U_{\lambda_0\lambda}\mathfrak{N}_{\lambda_0} = \mathfrak{N}_\lambda$. To prove (4.10) we assume that there is a function $f \in \mathcal{H}$ such that

$$f \perp \text{c.l.s.} \quad \mathfrak{N}_\lambda.$$

$\lambda \neq \bar{\lambda}, \lambda \in \rho(T)$

Then $(f, U_{\lambda_0\lambda}g) = 0$ for all $g \in \mathfrak{N}_{\lambda_0}$ and all $\lambda \in \rho(T)$. But accretiveness of the system Θ implies that there are regular points of T in the upper and lower half-planes. This leads to a conclusion that the function $\phi(\lambda) = (f, U_{\lambda_0\lambda}g) \equiv 0$ for all $\lambda \neq \bar{\lambda}$. Combining this with (3.6) we conclude that $f = 0$ and thus (4.10) holds. □

5. RESTORING A NON-SELF-ADJOINT SCHRÖDINGER OPERATOR T_h

In this section we are going to use the realization results for Stieltjes functions developed in section 4 to obtain the solution of inverse spectral problem for Schrödinger operator of the form (2.5) in $L_2[a, +\infty)$ with non-self-adjoint boundary conditions

$$(5.1) \quad \begin{cases} T_h y = -y'' + q(x)y \\ y'(a) = hy(a) \end{cases}, \quad \left(q(x) = \overline{q(x)}, \operatorname{Im} h \neq 0 \right).$$

In particular, we will show that if a non-decreasing function $\sigma(t)$ is the spectral function of positive self-adjoint boundary value problem

$$(5.2) \quad \begin{cases} A_\theta y = -y'' + q(x)y \\ y'(a) = \theta y(a) \end{cases}$$

and satisfies conditions

$$(5.3) \quad \int_0^\infty d\sigma(t) = \infty, \quad \int_0^\infty \frac{d\sigma(t)}{1+t} < \infty,$$

then for every $\gamma \geq 0$ a Stieltjes function

$$V(z) = \gamma + \int_0^\infty \frac{d\sigma(t)}{t-z}$$

can be realized in the unique way as a $V_\Theta(z)$ function of an accretive rigged canonical system Θ with some Schrödinger operator T_h .

Let $\mathcal{H} = L_2[a, +\infty)$ and $l(y) = -y'' + q(x)y$ where q is a real locally summable function. We consider a symmetric operator

$$(5.4) \quad \begin{cases} \tilde{B}y = -y'' + q(x)y \\ y'(a) = y(a) = 0 \end{cases}$$

together with its positive self-adjoint extension of the form

$$(5.5) \quad \begin{cases} \tilde{B}_\theta y = -y'' + q(x)y \\ y'(a) = \theta y(a) \end{cases}$$

defined in $\mathcal{H} = L_2[a, +\infty)$. A non-decreasing function $\sigma(\lambda)$ defined on $[0, +\infty)$ is called the *distribution function* (see [23]) of an operator pair $\tilde{B}_\theta, \tilde{B}$, where $\tilde{B}_\theta$ of the form (5.5) is a self-adjoint extension of symmetric operator $\tilde{B}$ of the form (5.4), and if the formulas

$$(5.6) \quad \begin{aligned} \varphi(\lambda) &= Uf(x), \\ f(x) &= U^{-1}\varphi(\lambda), \end{aligned}$$

establish one-to-one isometric correspondence U between $L_2^\sigma[0, +\infty)$ and $L_2[a, +\infty)$. Moreover, this correspondence is such that the operator $\tilde{B}_\theta$ is unitarily equivalent to the operator

$$(5.7) \quad \Lambda_\sigma \varphi(\lambda) = \lambda \varphi(\lambda), \quad (\varphi(\lambda) \in L_2^\sigma[0, +\infty))$$

in $L_2^\sigma[0, +\infty)$ while symmetric operator $\tilde{B}$ in (5.4) is unitarily equivalent to the symmetric operator

$$(5.8) \quad \Lambda_\sigma^0 \varphi(\lambda) = \lambda \varphi(\lambda), \quad \left(\varphi(\lambda) \in L_2^\sigma[0, +\infty), \int_0^{+\infty} \varphi(\lambda) d\sigma(\lambda) = 0 \right).$$

Definition 5.1. A scalar Herglotz-Nevanlinna function $V(z)$ is called *Stieltjes like function* if it has an integral representation (4.1) with an arbitrary (not necessarily non-negative) constant γ .

We are going to introduce a new class of realizable scalar Stieltjes like functions whose structure is similar to that of $SL_0(R)$ of section 4.

Definition 5.2. A Stieltjes like function $V(z)$ is said to be a member of the *class* $SL_0(R)$ if it admits an integral representation

$$(5.9) \quad V(z) = \gamma + \int_0^\infty \frac{d\sigma(t)}{t-z}, \quad (\gamma \in (-\infty, +\infty)),$$

where non-decreasing function $\sigma(t)$ satisfies the following conditions

$$(5.10) \quad \int_0^\infty d\sigma(t) = \infty, \quad \int_0^\infty \frac{d\sigma(t)}{1+t} < \infty.$$

Consider the following subclasses of $SL_0(R)$.

Definition 5.3. A function $V(z) \in SL_0(R)$ belongs to the *class* $SL_0^K(R)$ if

$$(5.11) \quad \int_0^\infty \frac{d\sigma(t)}{t} = \infty.$$

Definition 5.4. A function $V(z) \in SL_0(R)$ belongs to the *class* $SL_{01}^K(R)$ if

$$(5.12) \quad \int_0^\infty \frac{d\sigma(t)}{t} < \infty.$$

The following theorem describes the realization of the class $SL_0(R)$.

Theorem 5.5. Let $V(z) \in SL_0(R)$ and the function $\sigma(t)$ be the distribution function of an operator pair $\tilde{B}_\theta$ of the form (5.4) and $\tilde{B}$ of the form (5.5). Then there exist unique Schrödinger operator T_h ($\text{Im } h > 0$) of the form (5.1), operator $\mathbb{A}$ given by (2.6), operator K as in (3.14), and the rigged canonical system of the Livsic type

$$(5.13) \quad \Theta = \begin{pmatrix} \mathbb{A} & K & 1 \\ \mathcal{H}_+ \subset L_2[a, +\infty) \subset \mathcal{H}_- & \mathbb{C} & \end{pmatrix},$$

of the form (3.15) so that $V(z)$ is realized by Θ .

Proof. Since $\sigma(t)$ is the distribution function of the positive self-adjoint operator, then (see [23]) we can completely restore the operator $\tilde{B}_\theta$ of the form (5.5) as well as a symmetric operator $\tilde{B}$ of the form (5.4). It follows from the definition of the distribution function above that there is operator U defined in (5.6) establishing one-to-one isometric correspondence between $L_2^\sigma[0, +\infty)$ and $L_2[a, +\infty)$ while providing for the unitary equivalence between the operator $\tilde{B}_\theta$ and operator of multiplication by independent variable Λ_σ of the form (5.7). Taking this into account, we realize (see [11]) a Herglotz-Nevanlinna function $V(z)$ with a rigged canonical system

$$\Theta_\Lambda = \begin{pmatrix} \Lambda & K^\sigma & 1 \\ \mathcal{H}_+^\sigma \subset L_2^\sigma[0, +\infty) \subset \mathcal{H}_-^\sigma & \mathbb{C} & \end{pmatrix}.$$

Following the steps for construction of the model system described in [11], we note that

$$\Lambda = \text{Re } \Lambda + iK^\sigma(K^\sigma)^*$$

is a correct $(*)$ -extension of an operator T^σ such that $\mathbf{\Lambda} \supset T^\sigma \supset \Lambda_\sigma^0$ where Λ_σ^0 is defined in (5.8). The real part $\text{Re } \mathbf{\Lambda}$ is a self-adjoint bi-extension of Λ_σ^0 that has a quasi-kernel Λ_σ of the form (5.7). The operator K^σ in the above system is defined by

$$K^\sigma c = c \cdot 1, \quad c \in \mathbb{C}, \quad 1 \in \mathcal{H}_-^\sigma.$$

Here we need to clarify why number 1 belongs to $\mathcal{H}_-^\sigma$. To confirm this we need to show that $(x, 1)$ defines a continuous linear functional for every $x \in \mathcal{H}_+^\sigma$. It was shown in [11], [12] that

$$\mathcal{H}_+^\sigma = \mathcal{D}(\Lambda_\sigma^0) \dot{+} \left\{ \frac{c_1}{1+t^2} \right\} \dot{+} \left\{ \frac{c_2 t}{1+t^2} \right\}, \quad c_1, c_2 \in \mathbb{C}.$$

Consequently, every vector $x \in \mathcal{H}_+^\sigma$ has three components $x = x_1 + x_2 + x_3$ according to the decomposition above. Obviously, $(x_1, 1)$ and $(x_2, 1)$ yield convergent integrals while $(x_3, 1)$ boils down to

$$\int_0^\infty \frac{t}{1+t^2} d\sigma(t).$$

To see the convergence of the above integral we notice that

$$\frac{t}{1+t^2} = \frac{t-1}{(1+t^2)(t+1)} + \frac{1}{1+t} \leq \frac{1}{1+t^2} + \frac{1}{1+t}.$$

The integrals taken of the last two expressions on the right side converge due to (1.2) and (5.10), and hence so does the integral of the left side. Thus, $(x, 1)$ defines a continuous linear functional for every $x \in \mathcal{H}_+^\sigma$ and $1 \in \mathcal{H}_-^\sigma$.

The state space of the system Θ_Λ is $\mathcal{H}_+^\sigma \subset L_2^\sigma[0, +\infty) \subset \mathcal{H}_-^\sigma$, where $\mathcal{H}_+^\sigma = \mathcal{D}((\Lambda_\sigma^0)^*)$. By the realization theorem [11] we have that $V(z) = V_{\Theta_\Lambda}(z)$.

We can also show that the system Θ_Λ is a prime system. In order to do so we need to show that

$$(5.14) \quad \text{c.l.s.}_{\lambda \neq \bar{\lambda}, \lambda \in \rho(T^\sigma)} \mathfrak{N}_\lambda = L_2^\sigma[0, +\infty),$$

where $\mathfrak{N}_\lambda$ are defect subspaces of the symmetric operator Λ_σ^0 . It is known (see [11]) that Λ_σ^0 is a prime operator. Hence we can follow the reasoning of the proof of theorem 4.5 and only confirm that operator T^σ has regular points in the upper and lower half-planes. To see this we first note that non-negative operator Λ_σ^0 has no kernel spectrum [1] on the left real half-axis. Then we apply Theorem 1 of [1] (see page 149 of vol. 2 of [1]) that gives the complete description of the spectrum of T^σ . This theorem implies that there are regular points of T^σ on the left real half-axis. Since $\rho(T^\sigma)$ is an open set we confirm the presence of non-real regular points of T^σ in both half-planes. Thus (5.14) holds and Θ_Λ is a prime system.

Applying theorem 3.2 on unitary equivalence to the isometry U defined in (5.6) we obtain a triplet of isometric operators U_+ , U , and U_- , where

$$U_+ = U|_{\mathcal{H}_+^\sigma}, \quad U_-^* = U_+^*.$$

This triplet of isometric operators will map the rigged Hilbert space Θ_Λ is $\mathcal{H}_+^\sigma \subset L_2^\sigma[0, +\infty) \subset \mathcal{H}_-^\sigma$ into another rigged Hilbert space $\mathcal{H}_+ \subset L_2^\sigma[a, +\infty) \subset \mathcal{H}_-$. Moreover, U_+ is an isometry from $\mathcal{H}_+^\sigma = \mathcal{D}(\Lambda_\sigma^{0*})$ onto $\mathcal{H}_+ = \mathcal{D}(\tilde{B}^*)$, and $U_-^* = U_+^*$ is an isometry from $\mathcal{H}_+^\sigma$ onto $\mathcal{H}_-$. This is true since the operator U provides the unitary equivalence between the symmetric operators $\tilde{B}$ and Λ_σ^0 .

Now we construct a system

$$\Theta = \begin{pmatrix} \mathbb{A} & K & 1 \\ \mathcal{H}_+ \subset L_2[a, +\infty) \subset \mathcal{H}_- & \mathbb{C} & \end{pmatrix}$$

where $K = U_- K^\sigma$ and $\mathbb{A} = U_- \Lambda U_+^{-1}$ is a correct $(*)$ -extension of operator $T = UT^\sigma U^{-1}$ such that $\mathbb{A} \supset T \supset \tilde{B}$. The real part $\operatorname{Re} \mathbb{A}$ contains the quasi-kernel $\tilde{B}_\theta$. This construction of $\mathbb{A}$ is unique due to the theorem on the uniqueness of a $(*)$ -extension for a given quasi-kernel (see [27]). On the other hand, all $(*)$ -extensions based on a pair $\tilde{B}, \tilde{B}_\theta$ must take form (2.6) for some values of parameters h and μ . Consequently, our function $V(z)$ is realized by the system Θ of the form (5.13) and

$$V(z) = V_{\Theta_\Lambda}(z) = V_\Theta(z).$$

□

Remark 5.6. Applying corollary 3.3 to the mapping U defined by (5.6) we obtain that the operator U in the above theorem is unique. The uniqueness of the operator U leads to an interesting observation. Let $u_k(x, \lambda)$, $(k = 1, 2)$ be the solutions of the following Cauchy problems:

$$\begin{cases} l(u_1) = \lambda u_1 \\ u_1(a, \lambda) = 0 \\ u_1'(a, \lambda) = 1 \end{cases}, \quad \begin{cases} l(u_2) = \lambda u_2 \\ u_2(a, \lambda) = 1 \\ u_2'(a, \lambda) = 0 \end{cases}.$$

Traditionally, (see [23]) a non-decreasing function $\sigma(\lambda)$ defined on $[0, +\infty)$ is called the *distribution function of a self-adjoint operator* $\tilde{B}_\theta$ of the form (5.5) if the formulas

$$(5.15) \quad \begin{aligned} \varphi(\lambda) &= U f(x) = \int_a^{+\infty} f(x) u(x, \lambda) dx, \\ f(x) &= U^{-1} \varphi(\lambda) = \int_0^{+\infty} \varphi(\lambda) u(x, \lambda) d\sigma(\lambda), \end{aligned}$$

where $u(x, \lambda) = u_1(x, \lambda) + \theta u_2(x, \lambda)$, establish one-to-one isometric correspondence U between $L_2^\sigma[0, +\infty)$ and $L_2[a, +\infty)$ such that the operator $\tilde{B}_\theta$ in (5.5) is unitarily equivalent to the operator Λ_σ in (5.7). It is easily seen that if the mapping U in (5.15) is such that symmetric operators $\tilde{B}$ in (5.4) and Λ_σ^0 in (5.8) are unitarily equivalent w.r.t. U as well, then the mapping U in theorem 5.5 is given by the formulas (5.15). Indeed, assuming that there is another mapping $\tilde{U}$ provided by theorem 3.2 on unitary equivalence for the systems Θ_Λ and Θ we would violate the uniqueness of the operator U , and thus $\tilde{U} = U$.

Theorem 5.7. *Let $V(z) \in SL_0(R)$ satisfy the conditions of theorem 5.5. Then the operator T_h in the conclusion of the theorem 5.5 is accretive if and only if*

$$(5.16) \quad \gamma^2 + \gamma \int_0^\infty \frac{d\sigma(t)}{t} + 1 \geq 0.$$

The operator T_h is α -sectorial for some $\alpha \in (0, \pi/2)$ if and only if the inequality (5.16) is strict. In this case the exact value of angle α can be calculated by the formula

$$(5.17) \quad \tan \alpha = \frac{\int_0^\infty \frac{d\sigma(t)}{t}}{\gamma^2 + \gamma \int_0^\infty \frac{d\sigma(t)}{t} + 1}.$$

Proof. It was shown in [26] that for the system Θ in (5.13) described in the previous theorem the operator T_h is accretive if and only if the function

$$(5.18) \quad \begin{aligned} V_h(z) &= -i[W_\Theta^{-1}(-1)W_\Theta(z) + I]^{-1}[W_\Theta(-1)W_\Theta(z) - I] \\ &= -i \frac{1 - [(m_\infty(z) + \bar{h})/(m_\infty(z) + h)][(m_\infty(-1) + h)/(m_\infty(-1) + \bar{h})]}{1 + [(m_\infty(z) + \bar{h})/(m_\infty(z) + h)][(m_\infty(-1) + h)/(m_\infty(-1) + \bar{h})]}, \end{aligned}$$

is holomorphic in $\text{Ext}[0, +\infty)$ and satisfies the following inequality

$$(5.19) \quad 1 + V_h(0) V_h(-\infty) \geq 0.$$

Here $W_\Theta(z)$ is the transfer function of (5.13). It is also shown in [26] that the operator T_h is α -sectorial for some $\alpha \in (0, \pi/2)$ if and only if the inequality (5.19) is strict while the exact value of angle α can be calculated by the formula

$$(5.20) \quad \cot \alpha = \frac{1 + V_\Theta(0) V_\Theta(-\infty)}{|V_\Theta(-\infty) - V_\Theta(0)|}.$$

According to theorem 5.5 and equation (3.5)

$$W_\Theta(z) = (I - iV(z)J)(I + iV(z)J)^{-1}.$$

By direct calculations one obtains

$$(5.21) \quad W_\Theta(-1) = \frac{1 - i \left[\gamma + \int_0^\infty \frac{d\sigma(t)}{t+1} \right]}{1 + i \left[\gamma + \int_0^\infty \frac{d\sigma(t)}{t+1} \right]}, \quad W_\Theta^{-1}(-1) = \frac{1 + i \left[\gamma + \int_0^\infty \frac{d\sigma(t)}{t+1} \right]}{1 - i \left[\gamma + \int_0^\infty \frac{d\sigma(t)}{t+1} \right]}.$$

Using the following notations

$$a = \int_0^\infty \frac{d\sigma(t)}{t+1} \quad \text{and} \quad b = \int_0^\infty \frac{d\sigma(t)}{t},$$

and performing straightforward calculations we obtain

$$(5.22) \quad V_h(0) = \frac{a - b}{1 + ab} \quad \text{and} \quad V_h(-\infty) = \frac{a - \gamma}{1 + a\gamma}.$$

Substituting (5.22) into (5.20) and performing the necessary steps we get

$$(5.23) \quad \cot \alpha = \frac{1 + b\gamma}{b - \gamma} = \frac{\gamma^2 + \gamma \int_0^\infty \frac{d\sigma(t)}{t} + 1}{\int_0^\infty \frac{d\sigma(t)}{t}}.$$

Taking into account that $b - \gamma > 0$ we combine (5.19), (5.20) with (5.23) and this completes the proof of the theorem. $\square$

Corollary 5.8. *Let $V(z) \in SL_0(R)$ satisfy the conditions of theorem 5.5. Then the operator T_h in the conclusion of theorem 5.5 is accretive if and only if*

$$(5.24) \quad 1 + V(0) V(-\infty) \geq 0.$$

The operator T_h is α -sectorial for some $\alpha \in (0, \pi/2)$ if and only if the inequality (5.24) is strict. In this case the exact value of angle α can be calculated by the formula

$$(5.25) \quad \tan \alpha = \frac{V(-\infty) - V(0)}{1 + V(0) V(-\infty)}.$$

Proof. Taking into account that

$$V(0) = \gamma + \int_0^\infty \frac{d\sigma(t)}{t},$$

$V(z) = V_\Theta(z)$, and $V_\Theta(-\infty) = \gamma$, we use (5.16) and (5.17) to obtain (5.24) and (5.25). $\square$

Theorem 5.9. *Let $V(z) \in S_0(R)$ and satisfy the conditions of theorem 5.5. Then the system Θ of the form (5.13) is accretive and its symmetric operator A of the form (2.4) is such that its Kreĭn-von Neumann extension A_K of the form (2.8) does not coincide with its Friedrichs extension A_F of the form (2.9).*

Proof. The proof of the fact that Θ is accretive directly follows from the theorems 4.2 and 5.5. The second part follows from the theorem in [25] that states that a positive symmetric operator A admits a non-self-adjoint accretive extension T if and only if A_F and A_K do not coincide. $\square$

Below we will derive the formulas for calculation of the boundary parameter h in the restored Schrödinger operator T_h of the form (5.1). We consider two major cases.

Case 1. In the first case we assume that $\int_0^\infty \frac{d\sigma(t)}{t} < \infty$. This means that our function $V(z)$ belongs to the class $SL_{01}^K(R)$. In what follows we denote

$$b = \int_0^\infty \frac{d\sigma(t)}{t} \quad \text{and} \quad m = m_\infty(-0).$$

Suppose that $b \geq 2$. Then the quadratic inequality (5.16) implies that for all γ such that

$$(5.26) \quad \gamma \in \left(-\infty, \frac{-b - \sqrt{b^2 - 4}}{2}\right] \cup \left[\frac{-b + \sqrt{b^2 - 4}}{2}, +\infty\right)$$

the restored operator T_h is accretive. Clearly, this operator is extremal accretive if

$$\gamma = \frac{-b \pm \sqrt{b^2 - 4}}{2}.$$

In particular if $b = 2$ then $\gamma = -1$ and the function

$$V(z) = -1 + \int_0^\infty \frac{d\sigma(t)}{t - z}$$

is realized using an extremal accretive T_h .

Now suppose that $0 < b < 2$. For every $\gamma \in (-\infty, +\infty)$ the restored operator T_h will be accretive and α -sectorial for some $\alpha \in (0, \pi/2)$. Consider a function $V(z)$ defined by (5.9). Conducting realizations of $V(z)$ by operators T_h for different values of $\gamma \in (-\infty, +\infty)$ we notice that the operator T_h with the largest angle of sectoriality occurs when

$$(5.27) \quad \gamma = -\frac{b}{2},$$

and is found according to the formula

$$(5.28) \quad \alpha = \arctan \frac{b}{1 - b^2/4}.$$

This follows from the formula (5.17), the fact that $\gamma^2 + \gamma b + 1 > 0$ for all γ , and the formula

$$\gamma^2 + \gamma b + 1 = \left(\gamma + \frac{b}{2}\right)^2 + \left(1 - \frac{b^2}{4}\right).$$

Now we will focus on the description of the parameter h in the restored operator T_h .

It was shown in [6] that the quasi-kernel $\hat{A}$ of the realizing system Θ from theorem 5.5 takes a form

$$(5.29) \quad \begin{cases} \hat{A}y = -y'' + qy \\ y'(a) = \eta y(a) \end{cases}, \quad \eta = \frac{\mu \operatorname{Re} h - |h|^2}{\mu - \operatorname{Re} h}$$

On the other hand, since $\sigma(t)$ is also the distribution function of the positive self-adjoint operator, we can conclude that $\hat{A}$ equals to the operator $\tilde{B}_\theta$ of the form (5.5). This connection allows us to obtain

$$(5.30) \quad \theta = \eta = \frac{\mu \operatorname{Re} h - |h|^2}{\mu - \operatorname{Re} h}.$$

Assuming that

$$h = x + iy$$

we will use (5.30) to derive the formulas for x and y in terms of γ . First, to eliminate parameter μ , we notice that (3.16) and (3.5) imply

$$W_\Theta(\lambda) = \frac{\mu - h}{\mu - \bar{h}} \frac{m_\infty(\lambda) + \bar{h}}{m_\infty(\lambda) + h} = \frac{1 - iV(z)}{1 + iV(z)}.$$

Passing to the limit when $z \rightarrow -\infty$ and taking into account that $V(-\infty) = \gamma$ we obtain

$$\frac{\mu - h}{\mu - \bar{h}} = \frac{1 - i\gamma}{1 + i\gamma}.$$

Let us denote

$$(5.31) \quad a = \frac{1 - i\gamma}{1 + i\gamma}.$$

Solving (5.31) for μ yields

$$\mu = \frac{h - a\bar{h}}{1 - a}.$$

Substituting this value into (5.30) after simplification produces

$$\frac{x + iy - a(x - iy)x - (x^2 + y^2)(1 - a)}{x + iy - a(x - iy) - x(1 - a)} = \theta.$$

After straightforward calculations targeting to represent numerator and denominator of the last equation in standard form one obtains the following relation



$$(5.32) \quad x - \gamma y = \theta.$$

It was shown in [26] that the α -sectoriality of the operator T_h and (5.20) lead to

$$(5.33) \quad \tan \alpha = \frac{\operatorname{Im} h}{\operatorname{Re} h + m_\infty(-0)} = \frac{y}{x + m_\infty(-0)}.$$

Combining (5.32) and (5.33) one obtains

$$x - \gamma(x \tan \alpha + m_\infty(-0) \tan \alpha) = \theta,$$

FIGURE 1. $b > 2$ FIGURE 2. γ interval

or

$$x = \frac{\theta + \gamma m_{\infty}(-0) \tan \alpha}{1 - \gamma \tan \alpha}.$$

But $\tan \alpha$ is also determined by (5.17). Direct substitution of

$$\tan \alpha = \frac{b}{1 + \gamma(\gamma + b)}$$

into the above equation yields

$$x = \theta + \frac{[\theta + m_{\infty}(-0)]b\gamma}{1 + \gamma^2}.$$

Using the short notation and finalizing calculations we get

$$(5.34) \quad h = x + iy, \quad x = \theta + \frac{\gamma[\theta + m]b}{1 + \gamma^2}, \quad y = \frac{[\theta + m]b}{1 + \gamma^2}.$$

At this point we can use (5.34) to provide analytical and graphical interpretation of the parameter h in the restored operator T_h . Let



$$c = (\theta + m)b.$$

Again we consider three subcases.

Subcase 1: $b > 2$ Using basic algebra we transform (5.34) into

$$(5.35) \quad (x - \theta)^2 + \left(y - \frac{c}{2}\right)^2 = \frac{c^2}{4}.$$

Since in this case the parameter γ belongs to the interval in (5.26), we

FIGURE 3. $b < 2$ FIGURE 4. $b = 2$

can see that h traces the highlighted part of the circle on the figure 1 as γ moves from $-\infty$ towards $+\infty$. We also notice that the removed point $(\theta, 0)$ corresponds to the value of $\gamma = \pm\infty$ while the points h_1 and h_2 correspond to the values $\gamma_1 = \frac{-b-\sqrt{b^2-4}}{2}$ and $\gamma_2 = \frac{-b+\sqrt{b^2-4}}{2}$, respectively (see figure 2).

Subcase 2: $b < 2$ For every $\gamma \in (-\infty, +\infty)$ the restored operator T_h will be accretive and α -sectorial for some $\alpha \in (0, \pi/2)$. As we have mentioned above, the operator T_h achieves the largest angle of sectoriality when $\gamma = -\frac{b}{2}$. In this particular case (5.34) becomes

$$(5.36) \quad h = x + iy, \quad x = \frac{\theta(4 - b^2) - 2b^2m}{4 + b^2}, \quad y = \frac{4(\theta + m)b}{4 + b^2}.$$

The value of h from (5.36) is marked on the figure 3.

Subcase 3: $b = 2$ The behavior of parameter h in this case is depicted on the figure 4. It shows that in this case the function $V(z)$ can be realized using an extremal accretive T_h when $\gamma = -1$. The value of the parameter

h according to (5.34) then becomes

$$(5.37) \quad h = x + iy, \quad x = -m, \quad y = \theta + m.$$

Clockwise direction of the circle again corresponds to the change of γ from $-\infty$ to $+\infty$ and the marked value of h occurs when $\gamma = -1$.

Now we consider the second case.

Case 2. Here we assume that $\int_0^\infty \frac{d\sigma(t)}{t} = \infty$. This means that our function $V(z)$ belongs to the class $SL_0^K(R)$ and $b = \infty$. According to theorem 5.7 and formulas (5.16) and (5.17), the restored operator T_h is accretive if and only if

$$\gamma \geq 0,$$

and α -sectorial if and only if $\gamma > 0$. It directly follows from (5.17) that the exact value of the angle α is then found from

$$(5.38) \quad \tan \alpha = \frac{1}{\gamma}.$$

The latter implies that the restored operator T_h is extremal if $\gamma = 0$. This means that a function $V(z) \in SL_0^K(R)$ is realized by a system with an extremal operator T_h if and only if

$$(5.39) \quad V(z) = \int_0^\infty \frac{d\sigma(t)}{t - z}.$$

On the other hand since $\gamma \geq 0$ the function $V(z)$ is a Stieltjes function of the class $S_0(R)$. Applying realization theorems from [15] we conclude that $V(z)$ admits realization by an accretive system Θ of the form (3.1) with $\mathbb{A}_R$ containing the Krein-von Neumann extension A_K as a quasi-kernel. Here A_K is defined by (2.8). This yields

$$(5.40) \quad \theta = -m_\infty(-0) = -m.$$

As in the beginning of the previous case we derive the formulas for x and y , where $h = x + iy$. Using (5.30) and (5.32) leads to

$$(5.41) \quad \begin{cases} \theta = \frac{\mu x - (x^2 + y^2)}{\mu - x}, \\ x = \theta + \gamma y. \end{cases}$$

Solving this system for x and y leads to

$$(5.42) \quad x = \frac{\theta + \mu\gamma^2}{1 + \gamma^2}, \quad y = \frac{(\mu - \theta)\gamma}{1 + \gamma^2}.$$

Combining (5.41) and (5.42) gives



$$(5.43) \quad x = \frac{-m + \mu\gamma^2}{1 + \gamma^2}, \quad y = \frac{(m + \mu)\gamma}{1 + \gamma^2}.$$

To proceed, we first notice that our function $V(z)$ satisfies the conditions of theorem 4.8 of [6]. Indeed, the inequality

$$\mu \geq \frac{(\operatorname{Im} h)^2}{m_\infty(-0) + \operatorname{Re} h},$$

turns into

$$\mu = \frac{y^2}{x - m} + x,$$

FIGURE 5. $b = \infty$ FIGURE 6. $\gamma \geq 0$

if you use $\theta = -m$ and the first equation in (5.41). Applying theorem 4.8 of [6] yields

$$(5.44) \quad \int_0^\infty \frac{d\sigma(t)}{1+t^2} = \frac{\operatorname{Im} h}{|\mu - h|^2} \left(\sup_{y \in D(A_K)} \frac{|\mu y(a) - y'(a)|}{\left(\int_a^\infty (|y(x)|^2 + |l(y)|^2) dx \right)^{\frac{1}{2}}} \right)^2.$$

Taking into account that

$$\mu y(a) - y'(a) = (\mu + m)y(a)$$

and setting

$$(5.45) \quad c^{1/2} = \sup_{y \in D(A_K)} \frac{|y(a)|}{\left(\int_a^\infty (|y(x)|^2 + |l(y)|^2) dx \right)^{\frac{1}{2}}},$$

we obtain

$$(5.46) \quad \frac{\operatorname{Im} h}{|\mu - h|^2} (\mu + m)^2 c = \int_0^\infty \frac{d\sigma(t)}{1+t^2}.$$

Considering that $\operatorname{Im} h = y$ and combining (5.46) with (5.43) we use straightforward

calculations to get

$$\mu = -m + \left(\frac{1}{\gamma c} \right) \int_0^\infty \frac{d\sigma(t)}{1+t^2}.$$

Let

$$(5.47) \quad \xi = \frac{1}{c} \int_0^\infty \frac{d\sigma(t)}{1+t^2}.$$

Then the last equation becomes

$$(5.48) \quad \mu = -m + \frac{\xi}{\gamma}.$$

Applying (5.48) on (5.43) yields

$$(5.49) \quad x = -m + \frac{\gamma \xi}{1+\gamma^2}, \quad y = \frac{\xi}{1+\gamma^2}, \quad \gamma \geq 0.$$

Following the previous case approach we transform (5.49) into

$$(5.50) \quad (x+m)^2 + \left(y - \frac{\xi}{2} \right)^2 = \frac{\xi^2}{4}.$$

The connection between the parameters γ and h in the accretive restored operator T_h is depicted in figures 5 and 6. As we can see h traces the highlighted part of the circle clockwise on the figure 5 as γ moves from 0 towards $+\infty$.

As we mentioned earlier the restored operator T_h is extremal if $\gamma = 0$. In this case formulas (5.49) become

$$(5.51) \quad x = -m, \quad y = \xi, \quad \gamma = 0,$$

where ξ is defined by (5.47).

6. REALIZING SYSTEMS WITH SCHRÖDINGER OPERATOR

Now once we described all the possible outcomes for the restored accretive operator T_h , we can concentrate on the main operator $\mathbb{A}$ of the system (5.13). We recall that $\mathbb{A}$ is defined by formulas (2.6) and beside the parameter h above contains also parameter μ . We will obtain the behavior of μ in terms of the components of our function $V(z)$ the same way we treated the parameter h . As before we consider two major cases dividing them into subcases when necessary.

Case 1. Assume that $b = \int_0^\infty \frac{d\sigma(t)}{t} < \infty$. In this case our function $V(z)$ belongs to the class $SL_{01}^K(R)$. First we will obtain the representation of μ in terms of x and y , where $h = x + iy$. We recall that


$$\mu = \frac{h - a\bar{h}}{1 - a},$$

where a is defined by (5.31). By direct computations we derive that

$$a = \frac{1-\gamma^2}{1+\gamma^2} - \frac{2\gamma}{1+\gamma^2}i, \quad 1-a = \frac{2\gamma^2}{1+\gamma^2} + \frac{2\gamma}{1+\gamma^2}i,$$

and

$$h - a\bar{h} = \left(\frac{2\gamma^2}{1+\gamma^2}x + \frac{2\gamma}{1+\gamma^2}y \right) + \left(\frac{2}{1+\gamma^2}y + \frac{2\gamma}{1+\gamma^2}x \right)i.$$

FIGURE 7. $b > 2$

Plugging the last two equations into the formula for μ above and simplifying we obtain

$$(6.1) \quad \mu = x + \frac{1}{\gamma} y.$$

We recall that during the present case x and y parts of h are described by the formulas (5.34).

Once again we elaborate in three subcases.

Subcase 1: $b > 2$ As we have shown this above, the formulas (5.34) can be transformed into equation of the circle (5.35). In this case the parameter γ belongs to the interval in (5.26), the accretive operator T_h corresponds to the values of h shown in the bold part of the circle on the figure 1 as γ moves from $-\infty$ towards $+\infty$.

Substituting the expressions for x and y from (5.34) into (6.1) and simplifying we get


$$(6.2) \quad \mu = \theta + \frac{(\theta + m)b}{\gamma}.$$

The connection between values of γ and μ is depicted on the figure 7. We note that $\mu = 0$ when $\gamma = -\frac{(\theta+m)b}{\theta}$. Also, the endpoints

$$\gamma_1 = \frac{-b - \sqrt{b^2 - 4}}{2} \quad \text{and} \quad \gamma_2 = \frac{-b + \sqrt{b^2 - 4}}{2}$$

of γ -interval (5.26) are responsible for the μ -values

$$\mu_1 = \theta + \frac{(\theta + m)b}{\gamma_1} \quad \text{and} \quad \mu_2 = \theta + \frac{(\theta + m)b}{\gamma_2}.$$

FIGURE 8. $b < 2$ and $b = 2$

The values of μ that are acceptable parameters of operator $\mathbb{A}$ of the restored system make the bold part of the hyperbola on the figure 7. It follows from theorem 4.2 that the operator $\mathbb{A}$ of the form (2.6) is accretive if and only if $\gamma \geq 0$ and thus μ sweeps the right branch on the hyperbola. We note that figure 7 shows the case when $-m < 0$, $\theta > 0$, and $\theta > -m$. Other possible cases, such as $(-m < 0, \theta < 0, \theta > -m)$, $(-m < 0, \theta = 0)$, and $(m = 0, \theta > 0)$ require corresponding adjustments to the graph shown in the picture 7.

Subcase 2: $b < 2$ For every $\gamma \in (-\infty, +\infty)$ the restored operator T_h will be accretive and α -sectorial for some $\alpha \in (0, \pi/2)$. As we have mentioned above, the operator T_h achieves the largest angle of sectoriality when $\gamma = -\frac{b}{2}$. In this particular case (5.34) becomes

$$h = x + iy, \quad x = \frac{\theta(4 - b^2) - 2b^2m}{4 + b^2}, \quad y = \frac{4(\theta + m)b}{4 + b^2}.$$


Substituting $\gamma = b/2$ into (6.1) we obtain

$$(6.3) \quad \mu = -(\theta + 2m).$$

This value of μ from (6.3) is marked on the figure 8. The corresponding operator $\mathbb{A}$ of the realizing system is based on these values of parameters h and μ .

Subcase 3: $b = 2$ The behavior of parameter μ in this case is also shown on the figure 8. It was shown above that in this case the function $V(z)$ can be realized using an extremal accretive T_h when $\gamma = -1$. The values of the parameters h and μ then become

$$h = x + iy, \quad x = -m, \quad y = \theta + m, \quad \mu = -(\theta + 2m).$$

FIGURE 9. $b = \infty$

The value of μ above is marked on the left branch of the hyperbola and occurs when $\gamma = -1 = -b/2$.

Case 2. Again we assume that $\int_0^\infty \frac{d\sigma(t)}{t} = \infty$. Hence $V(z) \in SL_0^K(R)$ and $b = \infty$. As we mentioned above the restored operator T_h is accretive if and only if $\gamma \geq 0$ and α -sectorial if and only if $\gamma > 0$. It is extremal if $\gamma = 0$. The values of x , y , and μ were already calculated and are given in (5.49) and (5.48), respectively. That is

$$x = -m + \frac{\gamma \xi}{1 + \gamma^2}, \quad y = \frac{\xi}{1 + \gamma^2}, \quad \mu = -m + \frac{\xi}{\gamma}, \quad \gamma \geq 0.$$

where ξ is defined in (5.47). Figure 9 gives graphical representation of this case. Only the right bold branch of hyperbola shows the values of μ in the case $b = \infty$. If $m = 0$ then

$$\mu = \frac{\xi}{\gamma}$$

and the graph should be adjusted accordingly.

In the case when $\gamma = 0$ and T_h is extremal we have

$$(6.4) \quad x = -m, \quad y = \xi, \quad \mu = \infty, \quad h = -m + i\xi,$$

and according to (2.6) we have

$$(6.5) \quad \mathbb{A}y = -y'' + q(x)y + [(-m + i\xi)y(a) - y'(a)]\delta(x - a),$$

that is the main operator of the realizing system.

Example. We conclude this paper with simple illustration. Consider a function

$$(6.6) \quad V(z) = \frac{i}{\sqrt{z}}.$$

A direct check confirms that $V(z)$ is a Stieltjes function. It was shown in [23] (see pp. 140-142) that the inversion formula

$$(6.7) \quad \sigma(\lambda) = C + \lim_{y \rightarrow 0} \frac{1}{\pi} \int_0^\lambda \operatorname{Im} \left(\frac{i}{\sqrt{x + iy}} \right) dx$$

describes the distribution function for a self-adjoint operator

$$\begin{cases} \tilde{B}_0 y = -y'' \\ y'(0) = 0. \end{cases}$$

The corresponding to $\tilde{B}_0$ symmetric operator is

$$(6.8) \quad \begin{cases} B_0 y = -y'' \\ y(0) = y'(0) = 0. \end{cases}$$

It was also shown in [23] that $\sigma(\lambda) = 0$ for $\lambda \leq 0$ and

$$(6.9) \quad \sigma'(\lambda) = \frac{1}{\pi\sqrt{\lambda}} \quad \text{for } \lambda > 0.$$

By direct calculations one can confirm that

$$V(z) = \int_0^\infty \frac{d\sigma(t)}{t-z} = \frac{i}{\sqrt{z}},$$

and that

$$\int_0^\infty \frac{d\sigma(t)}{t} = \int_0^\infty \frac{dt}{\pi t^{3/2}} = \infty.$$

It is also clear that the constant term in the integral representation (4.1) is zero, i.e. $\gamma = 0$.

Let us assume that $\sigma(t)$ satisfies our definition of spectral distribution function of the pair $B_0, \tilde{B}_0$ given in the section 5. Operating under this assumption, we proceed to restore parameters h and μ and apply formulas (5.49) for the values $\gamma = 0$ and $m = -\theta = 0$. This yields $x = 0$. To obtain y we first find the value of

$$\int_0^\infty \frac{d\sigma(t)}{1+t^2} = \frac{1}{\sqrt{2}},$$

and then use formula (5.45) to get the value of c . This yields $c = 1/\sqrt{2}$. Consequently,

$$\xi = \frac{1}{c} \int_0^\infty \frac{d\sigma(t)}{1+t^2} = 1,$$

and hence $h = yi = i$. From (5.48) we have that $\mu = \infty$ and (6.5) becomes

$$(6.10) \quad \mathbb{A} y = -y'' + [iy(0) - y'(0)]\delta(x).$$

The operator T_h in this case is

$$\begin{cases} T_h y = -y'' \\ y'(0) = iy(0). \end{cases}$$

The channel vector g of the form (3.12) then equals

$$(6.11) \quad g = \delta(x),$$

satisfying

$$\text{Im } \mathbb{A} = \frac{\mathbb{A} - \mathbb{A}^*}{2i} = K K^* = (., g)g,$$

and channel operator $Kc = cg$, ($c \in \mathbb{C}$) with

$$(6.12) \quad K^* y = (y, g) = y(0).$$

The real part of $\mathbb{A}$

$$\text{Re } \mathbb{A} y = -y'' - y'(0)\delta(x)$$

contains the self-adjoint quasi-kernel

$$\begin{cases} \hat{A}y = -y'' \\ y'(0) = 0. \end{cases}$$

A system of the Livšic type with Schrödinger operator of the form (5.13) that realizes $V(z)$ can now be written as

$$\Theta = \begin{pmatrix} \mathbb{A} & K & 1 \\ \mathcal{H}_+ \subset L_2[a, +\infty) \subset \mathcal{H}_- & \mathbb{C} & \end{pmatrix}.$$

where $\mathbb{A}$ and K are defined above. Now we can back up our assumption on $\sigma(t)$ to be the spectral distribution function of the pair $B_0, \tilde{B}_0$. Indeed, calculating the function $V_\Theta(z)$ for the system Θ above directly via formula (3.17) with $\mu = \infty$ and comparing the result to $V(z)$ gives the exact value of $h = i$. Using the reasoning of remark 5.6 we confirm that $\sigma(t)$ is the spectral distribution function of the pair $B_0, \tilde{B}_0$.

Remark 6.1. All the derivations above can be repeated for a Stieltjes like function

$$V(z) = \gamma + \frac{i}{\sqrt{z}}, \quad -\infty < \gamma < +\infty, \quad \gamma \neq 0$$

with very minor changes. In this case the restored values for h and μ are described as follows:

$$h = x + iy, \quad x = \frac{\gamma}{1 + \gamma^2}, \quad y = \frac{1}{1 + \gamma^2}, \quad \mu = \frac{1}{\gamma}.$$

The dynamics of changing h according to changing γ is depicted on the figure 5 where the circle has the center at the point $i/2$ and radius of $1/2$. The behavior of μ is described by a hyperbola $\mu = 1/\gamma$ (see figure 9 with $\theta = 0$). In the case when $\gamma > 0$ our function becomes Stieltjes and the restored system Θ is accretive. The operators $\mathbb{A}$ and K of the restored system are given according to the formulas (2.6) and (3.14), respectively.

REFERENCES

- [1] N.I. Akhiezer and I.M. Glazman. *Theory of linear operators*. Pitman Advanced Publishing Program, 1981. [2](#), [2](#), [5](#)
- [2] D. Alpay, I. Gohberg, M. A. Kaashoek, A. L. Sakhnovich, “Direct and inverse scattering problem for canonical systems with a strictly pseudoexponential potential”, *Math. Nachr.* 215 (2000), 531. [1](#)
- [3] D. Alpay and E.R. Tsekanovskii, “Interpolation theory in sectorial Stieltjes classes and explicit system solutions”, *Lin. Alg. Appl.*, 314 (2000), 91–136. [1](#), [4](#)
- [4] Yu.M. Arlinskiĭ. *On regular (*)-extensions and characteristic matrix valued functions of ordinary differential operators*. Boundary value problems for differential operators, Kiev, 3–13, 1980. [2](#)
- [5] Yu. Arlinskiĭ and E. Tsekanovskii. *Regular (*)-extension of unbounded operators, characteristic operator-functions and realization problems of transfer functions of linear systems*. Preprint, VINITI, Dep.-2867, 72p., 1979. [2](#)
- [6] Yu.M. Arlinskiĭ and E.R. Tsekanovskii, “Linear systems with Schrödinger operators and their transfer functions”, *Oper. Theory Adv. Appl.*, 149, 2004, 47–77. [2](#), [2](#), [3](#), [3](#), [1](#), [3](#), [5](#), [5](#)
- [7] D. Arov, H. Dym, “Strongly regular J -inner matrix-valued functions and inverse problems for canonical systems”, *Oper. Theory Adv. Appl.*, 160, Birkhauser, Basel, (2005), 101–160. [1](#)

- [8] D. Arov, H. Dym, “Direct and inverse problems for differential systems connected with Dirac systems and related factorization problems”, *Indiana Univ. Math. J.* 54 (2005), no. 6, 1769–1815. [1](#)
- [9] J.A. Ball and O.J. Staffans, “Conservative state-space realizations of dissipative system behaviors”, *Integr. Equ. Oper. Theory* (Online), Birkhäuser, 2005, DOI 10.1007/s00020-003-1356-3. [1](#)
- [10] H. Bart, I. Gohberg, and M. A. Kaashoek, *Minimal Factorizations of Matrix and Operator Functions*, Operator Theory: Advances and Applications, Vol. 1, Birkhäuser, Basel, 1979.
- [11] S.V. Belyi and E.R. Tsekanovskii, “Realization theorems for operator-valued R -functions”, *Oper. Theory Adv. Appl.*, 98 (1997), 55–91. [1](#), [1](#), [1](#), [2](#), [4](#), [4](#), [4](#), [5](#), [5](#)
- [12] S.V. Belyi and E.R. Tsekanovskii, “On classes of realizable operator-valued R -functions”, *Oper. Theory Adv. Appl.*, 115 (2000), 85–112. [1](#), [1](#), [4](#), [4](#), [4](#), [5](#)
- [13] M.S. Brodskii, *Triangular and Jordan representations of linear operators*, Moscow, Nauka, 1969 (Russian). [1](#)
- [14] V.A. Derkach, E.R. Tsekanovskii, “On characteristic operator-functions of accretive operator colligations”, *Ukrainian Math. Dokl.*, vol. A, no. 8, (1981), 16–20 (Ukrainian). [4](#)
- [15] I. Dovzhenko and E.R. Tsekanovskii, “Classes of Stieltjes operator-functions and their conservative realizations”, *Dokl. Akad. Nauk SSSR*, 311 no. 1 (1990), 18–22. [1](#), [4](#), [4](#), [4](#), [4](#), [5](#)
- [16] F. Gesztesy and E.R. Tsekanovskii, “On matrix-valued Herglotz functions”, *Math. Nachr.*, 218 (2000), 61–138. [1](#)
- [17] F. Gesztesy, N.J. Kalton, K.A. Makarov, E. Tsekanovskii, “Some Applications of Operator-Valued Herglotz Functions”, *Operator Theory: Advances and Applications*, **123**, Birkhäuser, Basel, (2001), 271–321. [1](#)
- [18] S. Khrushchev, “Spectral Singularities of dissipative Schrödinger operator with rapidly decreasing potential”, *Indiana Univ. Math. J.*, 33 no. 4, (1984), 613–638. [1](#)
- [19] I. S. Kac and M. G. Krein, *R -functions—analytic functions mapping the upper halfplane into itself*, *Amer. Math. Soc. Transl. (2)* **103**, 1–18 (1974). [4](#), [4](#)
- [20] Kato T.: *Perturbation Theory for Linear Operators*, Springer-Verlag, 1966 [2](#), [5](#)
- [21] B. M. Levitan, *Inverse Sturm-Liouville Problems*, VNU Science Press, Utrecht, 1987.
- [22] M.S. Livšic, *Operators, oscillations, waves*, Moscow, Nauka, 1966 (Russian). [1](#)
- [23] M. A. Naimark, *Linear Differential Operators II*, F. Ungar Publ., New York, 1968. [5](#), [5](#), [5](#), [6](#), [6](#)
- [24] O.J. Staffans, “Passive and conservative continuous time impedance and scattering systems, Part I: Well posed systems”, *Math. Control Signals Systems*, 15, (2002), 291–315. [1](#)
- [25] E.R. Tsekanovskii, “Accretive extensions and problems on Stieltjes operator-valued functions realizations”, *Oper. Theory Adv. Appl.*, 59 (1992), 328–347. [2](#), [2](#), [3](#), [5](#)
- [26] E.R. Tsekanovskii, “Characteristic function and sectorial boundary value problems”, *Investigation on geometry and math. analysis*, Novosibirsk, **7**, (1987), 180–194. [3](#), [5](#), [5](#), [5](#)
- [27] E.R. Tsekanovskii and Yu.L. Shmul’yan, “The theory of bi-extensions of operators on rigged Hilbert spaces. Unbounded operator colligations and characteristic functions”, *Russ. Math. Surv.*, 32 (1977), 73–131. [2](#), [2](#), [2](#), [3](#), [3](#), [3](#), [5](#)
- [28] V.A. Yurko, *Inverse problems for differential operators*, Saratv State University Publ., Saratov, 1989 (in Russian). [1](#)

DEPARTMENT OF MATHEMATICS, TROY STATE UNIVERSITY, TROY, AL 36082, USA
E-mail address: sbelyi@trojan.troyst.edu

DEPARTMENT OF MATHEMATICS, NIAGARA UNIVERSITY, NY 14109, USA
E-mail address: tsekanov@niagara.edu